



Brian E. Whipker<sup>1</sup>  
bwhipker@ncsu.edu



W. Garrett Owen<sup>2</sup>  
wgowen@msu.edu

## Torenia: *Impatiens Necrotic Spot Virus (INSV)*

*Necrotic leaf spot and faint necrotic ringspots were observed on torenia. Further inspection revealed necrotic stem tissue. This Alert will aid in you identifying Impatiens necrotic spot virus (INSV) on torenia. A series of photos provided by this Alert will aid in you identifying this problem on torenia.*

Torenia (*Torenia fournieri*) are nice cascading plants that are excellent additions to combination pots or by themselves in hanging baskets. During a recent scouting of a greenhouse, a few scattered torenia plants were noticed to be slow growing and having a number of leaves exhibiting browning tissue (necrosis) (Fig. 1). A closer inspection revealed several leaves with extensive spotting (Fig. 2) and having larger areas of necrotic tissue (Fig. 3). Some flower buds also displayed necrosis (Fig. 4). With the waxy surface, it was difficult to notice, without some effort, a faint necrotic ring pattern on two of the leaves (Fig. 5). No leaf mottling was present.

In addition, necrosis can also develop on the stems. The necrotic canker will ultimately girdle the stem causing wilt. This is the stage

2017 Sponsors



Figure 1. Torenia plant with overall slow growth and leaf spotting.

<sup>1</sup> North Carolina State University

<sup>2</sup> Michigan State University

## e-GRO Alert

[www.e-gro.org](http://www.e-gro.org)

### CONTRIBUTORS

**Dr. Nora Catlin**

Floriculture Specialist  
Cornell Cooperative Extension - Suffolk County  
[nora.catlin@cornell.edu](mailto:nora.catlin@cornell.edu)

**Dr. Chris Currey**

Assistant Professor of Floriculture  
Iowa State University  
[ccurrey@iastate.edu](mailto:ccurrey@iastate.edu)

**Dr. Ryan Dickson**

Ext. Specialist for Greenhouse Management & Technologies  
University of New Hampshire  
[ryan.dickson@unh.edu](mailto:ryan.dickson@unh.edu)

**Thomas Ford**

Commercial Horticulture Educator  
Penn State Extension  
[tgf2@psu.edu](mailto:tgf2@psu.edu)

**Dan Gilrein**

Entomology Specialist  
Cornell Cooperative Extension - Suffolk County  
[dog1@cornell.edu](mailto:dog1@cornell.edu)

**Dr. Joyce Latimer**

Floriculture Extension & Research  
Virginia Tech  
[jlatime@vt.edu](mailto:jlatime@vt.edu)

**Dr. Roberto Lopez**

Floriculture Extension & Research  
Michigan State University  
[rglopez@msu.edu](mailto:rglopez@msu.edu)

**Dr. Neil Mattson**

Greenhouse Research & Extension  
Cornell University  
[neil.mattson@cornell.edu](mailto:neil.mattson@cornell.edu)

**Dr. Garrett Owen**

Floriculture Outreach Specialist - Michigan State Univ.  
[wgowen@msu.edu](mailto:wgowen@msu.edu)

**Dr. Rosa E. Raudales**

Greenhouse Extension Specialist  
University of Connecticut  
[rosa.raudales@uconn.edu](mailto:rosa.raudales@uconn.edu)

**Dr. Beth Scheckelhoff**

Ext. Educator – Greenhouse Systems  
The Ohio State University  
[scheckelhoff.11@osu.edu](mailto:scheckelhoff.11@osu.edu)

**Lee Stivers**

Extension Educator – Horticulture  
Penn State Extension, Washington County  
[ljs32@psu.edu](mailto:ljs32@psu.edu)

**Dr. Paul Thomas**

Floriculture Extension & Research  
University of Georgia  
[pthomas@uga.edu](mailto:pthomas@uga.edu)

**Dr. Ariana Torres-Bravo**

Horticulture/ Ag. Econ., Purdue University  
[torres2@purdue.edu](mailto:torres2@purdue.edu)

**Dr. Brian Whipker**

Floriculture Extension & Research - NC State Univ.  
[bwhipker@ncsu.edu](mailto:bwhipker@ncsu.edu)

**Heidi Wollaeger**

Floriculture Outreach Specialist - Michigan State Univ.  
[wollaega@anr.msu.edu](mailto:wollaega@anr.msu.edu)

Copyright © 2017

Where trade names, proprietary products, or specific equipment are listed, no discrimination is intended and no endorsement, guarantee or warranty is implied by the authors, universities or

most growers notice that there is a problem (Fig. 6) (see earlier Alert 2.04 with more stem necrosis photographs).

These are all typical signs of a virus infection of torenia. A plant was tested for impatiens necrotic spot virus (INSV). INSV was confirmed with an enzyme-linked immunosorbent assay (ELISA) test by Mike Munster of the NC State University Plant and Insect Clinic.

### Management

Once a plant has INSV or tomato spotted wilt virus (TSWV), the virus cannot be removed. Discarding infected plants is the only option, and this will help prevent the virus from spreading further. It is important to note that some plants may be asymptomatic, but still have INSV or TSWV. Since the primary method of spreading these viruses is via Western Flower thrips (*Frankliniella occidentalis*) feeding, it is critical to keep them under control. See e-GRO Alert 4.18 for management options.



Figure 2. Close-up of necrotic leaf spotting on torenia that is indicative of a virus infection.



Figure 3. Expanding necrotic areas on the leaf as a result of an impatiens necrotic spot virus (INSV) infection.

Cooperating Universities

**UConn**



Cornell University



The University of Georgia

IOWA STATE UNIVERSITY

**MICHIGAN STATE  
UNIVERSITY**

**NC STATE  
UNIVERSITY**



THE OHIO STATE  
UNIVERSITY

PENNSTATE



Cooperative Extension  
College of Agricultural Sciences

**PURDUE  
UNIVERSITY**



University of  
New Hampshire  
Cooperative Extension



**VirginiaTech**  
*Invent the Future®*

MAUMEE VALLEY GROWERS  
*Choose the Very Best.*



In cooperation with our local and  
state greenhouse organizations



Indiana  
FLOWER  
GROWERS  
Association



Michigan Floriculture Growers Council



CONNECTICUT  
GREENHOUSE  
GROWERS  
ASSOCIATION





Figure 4. Close-up view of necrosis of the flower bud due to an impatiens necrotic spot virus (INSV) infection.



Figure 5. Necrotic ringspot symptoms associated with impatiens necrotic spot virus (INSV) infection. The ringspot can be difficult to observe with waxy leafed plants (within the circle).



Figure 6. Necrotic stem tissue associated with an impatiens necrotic spot virus (INSV) infection.